

# THE EFFECT OF ALTERING THE BIOSTIMULANT APPLICATION RATE ON THE STEM HEIGHT OF REPLANTED OILSEED SUNFLOWER PLANTS

**Dilfuza Qoraboeva Jo‘rayevna**

Termez State University, Faculty of Natural Sciences, Termez, Uzbekistan.

**Abstract.** Microfertilizers, biostimulants, microfertilizers and growth substances contained in the immunostimulant have been found to have a positive effect on the growth of the local Dilbar variety, on the weight of 1000 seeds and on the yield. The use of MERS microfertilizer made it possible to increase the productivity of one plant by 362.9-430.4 grams, the use of Fitovak immunostimulant by 288.1-388.0 grams, and the use of Biodux biostimulant by 107.7-287.7 grams.

**Keywords:** biostimulant, sunflower, standard, immunostimulant, microfertilizer, norm, yield, growth, development.

**Introduction.** Biologically active substances (BAS), including plant phytohormones - regulators (stimulators) of plant growth and development (RPGD) are becoming increasingly important in modern conditions. Their application in agriculture, plant science and forestry has the potential to produce results that cannot be achieved by other means. The use of RPGD makes it possible to better realize the genetic potential of crops, to increase the resistance of plants to stress factors of biotic and abiotic nature, and as a result, to increase the yield and improve the quality. Plant growth stimulants are special nutrients that accelerate plant metabolism and stimulate the accumulation of green mass in representatives of the flora. Plant growth regulators

are now widely available. They are available in different types. Some of them affect the formation of roots, increase the germination rate of seeds, Others affect the development and growth of the stem more, regulate flowering, are able to regulate the formation of seeded nodes.

**The level of study of the problem.** One of the advanced technologies of sunflower cultivation is the use of mineral fertilizers, which is one of the main factors of increasing productivity. It is important to nutrition the sunflower in time according to its demand, therefore, it is possible to obtain a high-quality and high yield if the amount and duration of the mineral fertilizer applied to it are determined [2].

Plant growth regulators cannot replace mineral fertilizers, but complement them in the plant nutrition system, increase the coefficient of use of soil and fertilizers. "UZGUMI" biofertilizer is applied mainly before planting seeds and is sprayed when sunflowers form 3-5 leaves. In this case, the yield increases by 0.22-0.31 t/ha and the oil content by 0.3-0.5%. [7]

It is necessary to use the latest scientific achievements and apply the innovative technologies of our advanced farms in order to achieve high productivity in sunflower cultivation and obtain a large amount of income and profit from it [1].

Under the influence of plant regulators, it increases the effectiveness of pesticides applied to seeds and during the growing season. Some growth regulators

reduce white and gray rot diseases of sunflower, and sugar beet diseases by 30-50% or more [3,8].

Employees of the Institute of Agroresurc of Ukraine (Ponomarenko SP, 2003] note that for the agricultural producer growth regulators are no less valuable than mineral fertilizers and plant protection products. A new direction of improving sunflower production technology is the development of a system of effective use of modern plant growth regulators that control individual stages of plant growth and development to activate their immunity and, as a result, increase the yield and quality of sunflower seeds. (Sonin K.E., 2010; Petrichenko V.N., 2010). Therefore, the development of technology for the use of biostimulants that regulate growth and increase immunity is of the utmost importance. [4,5,6].

**Research results.** Several doses of microfertilizers and biostimulants were applied to the Dilbar variety of sunflower, and the effects on field germination of seeds, the transition of sunflower phases, the height of the plant, the number of leaves, the size of baskets, the number and weight of seeds were studied. According to experiments, the height of the stem of the local Dilbar variety of sunflower is about 2 meters because it is planted in irrigated land. In the experiment, the height of the Dilbar variety grew to an average height of 161.7 sm in the control variant, it was found that the use of biostimulants had a positive effect on the plants, and the height of the plant grew from an average of 179.0 sm to 213.8 sm in these variants.

When UzGUMI biostimulant was applied to sunflower seeds at 0.6 l/ton, the average height of the plant was 197.6 sm, and the biostimulant had a positive effect on the growth of the plant stem, making it 35.9 sm taller than the control option. In the variant where Fitovak immunostimulant was applied to the seed in the amount of 300 ml/ton, the height of the plant was found to be 213.8 sm, and it was 52.1 sm higher than the control and 16.2 sm higher than the standard UzGUMI. In the variant where Fitovak immunostimulant was used in the amount of 200 ml/ton per seed, it was observed that the height of the plant grew at the same height (213.4 sm) as in the variant where the quantity of 300 ml/ton was used in the seed, but increasing the rate of the immunostimulant to 400 ml/ton per seed showed a decrease in the effect on the growth of the plant stem, and the stem height increased by 15.2 sm compared to the second option and 14.8 sm compared to the first option. The use of Fitovak immunostimulant made it possible to increase the height of stems by 51.7 sm compared to the first option, 52.1 compared to the second option, and 36.9 sm compared to the third option compared to the control option.

**Discussion.** If increasing the rate of MERS microfertilizer led to the growth of plant stem, it was found that in Bioduks and Fitovak biostimulants, on the contrary, increasing the rate of biostimulants had a negative effect on the growth of plant height.

## REFERENCES

1. Варшавская В.Б. Стимулирование прорастания семян сахарной свёклы регуляторами роста и другими физиологически активными веществами / Физиология семян: формирование, прорастание, прикладные аспекты. – Душанбе, 1990. – С. 311–314.
2. Нурматов Ш.Н., Азизов Т.Б., Турсунов Л., Анарбоев И.У., ва бошқалар //Мойли экинлардан юқори ҳосил етиштириш агротехнологияси бўйича тавсиялар. Тошкент. 2012. Туро-Иқбол нашриёти – 56–57 б
3. Мельников Н.Н. Пестициды и регуляторы роста растений / М.: Химия, 1995. – 576 с.
4. Петриченко В.Н. Влияние регуляторов роста растений и микроэлементов на урожайность подсолнечника и масличность семян / Аграр. Россия. – 2010. – №4. – С. 24– 26.
5. Пономаренко С.П. Регуляторы роста растений // – К.: 2003. – 319 с.
6. Сонин К.Е. Влияние препарата фуролан на формирование качества семян трёх сортов подсолнечника // Пищевая технология. – 2010. – №1. – С. 13–15.
7. Yuldasheva Z. K., Karabaeva D. J The effect of a biostimulator on the growth, development and yield of oily sunflower /“International Journal on Integrated Education” 2020. 157-160
8. Yuldasheva Z. K., Karabaeva D. J. Effect of biostimulator on the vegetation period of oily sunflower / «International journal for innovative engineering and management research» 2020. 122-125.
9. Халиков Б.М., Абдуалимов Ш ва бошқалар “УЗГУМИ” биоўғитидан фойдаланиш бўйича деҳқон ва фермер хўжаликларига тавсиялар. Тошкент, 2013, 4 б.
10. Z F Yuldasheva, D J Karabayeva. The effect of different doses of

different biostimulants on the yield of oily sunflower// IOP Conference Series: Earth and Environmental Science- 1142(2023) 012097